



FACULDADE DE MEDICINA  
UNIVERSIDADE DO PORTO

## **MESTRADO INTEGRADO EM MEDICINA**

---

2012/2013

Ricardo Filipe Teixeira Barbosa

The use of shotguns in violent  
deaths in the northern Portugal,  
2005 - 2010

março, 2013

FMUP



FACULDADE DE MEDICINA  
UNIVERSIDADE DO PORTO

Ricardo Filipe Teixeira Barbosa  
The use of shotguns in violent deaths  
in the northern Portugal, 2005-2010

**Mestrado Integrado em Medicina**

**Área: Medicina Legal**

**Trabalho efetuado sob a Orientação de:  
Professor Doutor Agostinho José Carvalho dos Santos**

**Trabalho organizado de acordo com as normas da revista:  
International Journal of Legal Medicine**

março, 2013

FMUP

Eu, Ricardo Filipe Teixeira Barbosa, abaixo assinado, nº mecanográfico 200605693, estudante do 6º ano do Mestrado Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 27/03/2013

Assinatura' Ricardo Filipe Teixeira Barbosa

**Nome:** Ricardo Filipe Teixeira Barbosa

**Email:** med06115@med.up.pt

**Título da Dissertação:**

The use of shotguns in violent deaths in the northern Portugal, 2005 - 2010

**Orientador:**

Professor Doutor Agostinho José Carvalho dos Santos

**Ano de conclusão:** 2013

**Designação da área do projeto:**

Medicina Legal

É autorizada a reprodução integral desta Dissertação para efeitos de investigação e de divulgação pedagógica, em programas e projetos coordenados pela FMUP.

Faculdade de Medicina da Universidade do Porto, 27/03/2013

Assinatura: Ricardo Filipe Teixeira Barbosa

# The use of shotguns in violent deaths in the northern Portugal, 2005 - 2010

Ricardo Barbosa <sup>a</sup>, Agostinho Santos <sup>a,b</sup>

<sup>a</sup> Faculty of Medicine of Porto University

<sup>b</sup> National Institute of Legal Medicine and Forensic Sciences – North Branch

## Authors' contact:

Ricardo Filipe Teixeira Barbosa  
Faculty of Medicine of Porto University  
Alameda Prof. Hernâni Monteiro 4200 - 319 Porto, Portugal  
(+351) 919097269/med06115@med.up.pt

## Abstract

Shotguns are used in a significant number of homicides and suicides. For cultural reasons these weapons are very common in Portugal. The main objective of this work is to improve the knowledge about the forensic aspects of all cases involving shotguns.

We conducted a retrospective study by analyzing all forensic autopsy reports involving the use of shotguns performed between January 1st 2005 and December 31st 2010, at the Northern Services of the National Institute of Legal Medicine and Forensic Sciences (n=105).

We evaluate 105 cases, with 54 (51.4%) homicides, 41 (39.0%) suicides and 10 (9.5%) accidents. Fifty-eight percent of them were male and 66.7% occurred in rural areas. Suicides occurred at home in 90.2%, while 48.1% of homicides and 66.7% of accidents occurred in public places. Homicides were caused by problems in the marital relationship in 32.1%; 25.0% of suicides by psychiatric diseases and 40.0% of accidents by mishandling of the gun. "Near contact" (78.9%) was the most common firing distance in suicides and the head (75.6%) was the most common location of the entry wound. For homicides, the most common firing distance was distant (59.1%), and the most prevalent distributions of entry wounds were head, chest (29.6% each), and multiple wound locations (37.0%).

Violent deaths caused by shotguns were not so scarce. The knowledge of this information can induce a change in mentalities to a more restrictive law that could reduce the number of violent deaths by firearms.

**Keywords:** firearms, shotguns, violent deaths, forensic autopsy

## **Introduction**

The use of firearms in violent deaths has been increasing worldwide and it is estimated that hundreds of thousands of people die annually [1]. In addition to fatalities, many more suffer non-fatal injuries, with serious physical and psychological consequences [1]. These cases constitute a major public health problem in many countries [2]. In the USA, in 2009, about 50% of all violent deaths, concerning 8200 cases, were due to the use of firearms [3]. In Portugal, according to statistics from the Ministry of Justice, the use of firearms was involved in 1311 crimes, between 2005 and 2010 [4].

People with guns at home, regardless the type of weapon, have a significantly higher risk of committing homicide and suicide by firearms than people without it [5]. In Portugal, with a population around 10,500 million people, it is estimated that there are around 2.6 million firearms [6], with an estimated value of 25 guns per 100 inhabitants, i.e. a quarter of the Portuguese population [6,7]. It is anticipated that about half of these guns are illegal [7].

In the international literature there are several studies that seek to analyze and characterize the cases of violent deaths caused by firearms [8-15]. However, these studies perform a general analysis of weapons and there are few specific studies to each type of gun, usually shotguns [13,16]. These weapons are, in some countries, the method of choice used in violent deaths, as is the case of Australia, United Kingdom and Scandinavian countries [9].

A shotgun is a weapon that is intended to be fired from the shoulder; it has a smooth bore and is designed to fire multiple pellets from the barrel [17]. Several factors justify the use of shotguns such as geographical aspects, socio-cultural and legal norms of the country in question [9,16]. In USA, the total number of deaths caused by shotguns has been declining in recent years, mainly through the reduction of homicide cases [13]. Males are the main victims of these weapons [13]. Its use in suicides is uncommon and it appears to be more related to rural areas [2]. Contrary to suicide and homicide cases, its involvement in accidents generally exceeds the use of handguns, in which hunting and mishandling of the firearm are the main causes to justify these accidents [8,16].

For historical and cultural reasons shotguns emerge as a very common practice in Portugal [6]. This fact may be related to its easiest legal acquisition to practice sport or hunting [6]. Prevention associated with behavior changes as well as changes to current national norms are keys to reduce deaths from firearms [18].

Given this reality, the main objective of this work is to promote knowledge about the forensic aspects of all cases involving shotguns. Specific aims are to characterize the use of shotguns in violent deaths that occurred in the northern region of Portugal and compare their relative frequency of use with the different manners of death (homicide, suicide or accident). According to our knowledge it is the first time that this kind of study is performed in Portugal.

## **Materials and Methods**

We conducted a retrospective study by analyzing all forensic autopsy reports involving the use of shotguns performed between January 1st 2005 and December 31st 2010, at the Northern Services of the National Institute of Legal Medicine and Forensic Sciences (n=105).

The database was constructed and statistical analysis performed on "SPSS - Social Package for Social Sciences v20.0". The variables were analyzed comparing the cases of alleged homicide, suicide and accident according to social and demographic characteristics of the victims, knowledge of the circumstances preceding death and forensic findings.  $p$  values of  $< 0,05$  were considered statistically significant.

## Results

### 1. Global Analysis

Shotguns were used in 105 violent deaths in the northern Portugal. According to forensic autopsy reports, the use of shotguns was found in 51.4% ( $n = 54$ ) of alleged homicides, 39.0% ( $n = 41$ ) of suicides and 9.5% ( $n = 10$ ) of accidents.

#### 1.1 Victims' characterization

Of 105 cases analyzed, 81.0% ( $n = 85$ ) of the victims were male and 19.0% ( $n = 20$ ) were female. Most victims lived in rural areas (66.7%,  $n = 70$ ). The age range was 16–82 years-old with a homogeneous distribution between groups starting from age 25yo. Most of the victims were married ( $n = 66$ ; 63.5%) and employed ( $n = 46$ , 43.8%) (see table 1). The current or previous occupation was known in 64 cases: farmer ( $n = 16$ , 25.0%) and trader ( $n = 10$ , 15.6%) were the most frequent. Only 3 victims were in occupation which enabled access to firearms (military, police officer and security).

Information regarding the history of diseases had been noted in 45 victims (57.7%) in a total of 78 cases in which we had access to the information. Cardiovascular system diseases were more frequent (46.7%,  $n = 21$ ), followed by the psychiatric diseases (31.1%,  $n = 14$ ).

The information about alcohol consumption was obtained in 74 cases: 42 (56.8%) were current consumers at the time of death and 30 (40.5%) had no history of alcohol consumption.

#### 1.2 Characterization of the event

Table 2 shows the division of cases according to the year of study. It is noted that the number of cases is very similar throughout the years and it was not possible to evaluate a trend in the use of these weapons over time. October ( $n = 16$ , 15.2%), August ( $n = 14$ , 13.3%) and December ( $n = 11$ , 10.5%) were the months with more cases reported.

Most cases occurred at home ( $n = 66$ ; 56.7%) or in a public place ( $n = 20$ ; 34.6%) (see table 3). Medical intervention occurred in only 16 cases (15.2%) and of these 12 (75.0%) had a lifetime of less than 24 hours and 4 survived more than 24 hours (25.0%).

#### 1.3. Forensic findings

The main forensic medical findings are resumed in table 4. The number of entrance wounds was established in 94 cases and in the remaining 11 cases the gunshot caused a slough of the tissue, which prevented the entrance wound characterization.

Fatal wounds were observed most in the head and/or neck in 50 cases (47.6%) or in the chest in 22 cases (21.0%); 20 cases had fatal wounds in multiple locations (37.0%).

The toxicological study was performed in 84 cases (80.0%) and toxicological tests were positive for ethanol (0.13–3.45 g/L) in 29 of those cases (34.5%). The presence of therapeutic drugs in the blood was detected in 3 cases, but all cases had concentrations of drugs in therapeutic ranges.

## 2. Analysis according to the manner of death

### 2.1 Victims' characterization

Table 5 shows the main social and demographic characteristics of the victims according to the manner of death. Victims were male in all accidents, and in suicides we have only recorded one female victim aged 37. This difference was smaller in homicides (men 64.8%, women 35.2%).

Homicides and suicides occurred more in adults, the group [25-64yo] had 72.2% and 65.9% of cases, respectively. In accidents, 40.0% ( $n = 4$ ) of the victims are in the group of [15-24yo], which corresponds to 80.0% of all cases occurring in this age group.

In suicides and accidents, farmer was the main occupation found, but note that accidents occurred in 3 students (30.0%). In homicides housewife was the occupation recorded in 20.6% ( $n = 6$ ) of cases, which corresponds to 66.6% of women in which the current or previous occupation was known.

Psychiatric diseases had been noted in 13 cases (31.7%) of all suicides, which correspond to 92.9% of all victims with psychiatric diseases. Depression was the main problem found.

### 2.2 Characterization of the event

If we analyze separately, June ( $n = 7$ , 17.1%) was the month in which there were more suicides, homicides occurred more in August ( $n = 10$ , 18.5%) and October ( $n = 5$ , 50.0%) was the month in which there were more accidents. The distribution according to different years of study was uniform.

We emphasize that almost all suicides occurred at home ( $n = 35$ , 90.2%), while homicides and accidents occurred mainly in public places ( $n = 26$ , 48.1% and  $n = 6$ , 66.7%, respectively) (see table 6).

According to the manner of death the table 7 shows motives/precipitating factors that have led to the event. In homicides, the problems in the marital relationship ( $n = 9$ , 32.1%), discussions with neighbors ( $n = 7$ , 25.0%) and robberies ( $n = 6$ , 21.4%) were the main causes found. In suicides, psychiatric diseases ( $n = 5$ , 25.0%) and life-threatening illness ( $n = 4$ , 9.8%) emerged as the main reasons. With regard to accidents, they occurred mostly by mishandling of the gun ( $n = 6$ , 60.0%) and hunting accidents ( $n = 3$ , 30.0%).

### 2.3. Forensic findings

Table 8 shows the main forensic findings according to the manner of death. One entrance wound was founded in all suicides. Head and/or neck ( $n = 31$ , 75.6%) were the most common anatomical regions, followed by chest and abdomen with 5 cases each (12.2%). In this situation the firing distance was known in 19 cases (46.3%): 15 (78.9%) were at “near contact” and 4 (21.1%) in contact.



With regard to homicides in 26 cases (48.1%) multiple entrance wounds (4 or more) were observed and in 22 cases (40.7%) the presence of only one entrance wound was reported. When only one anatomical region was involved head and/or neck and chest were the most affected with 16 cases each (29.6%). Note that in homicides the simultaneous achievement of various anatomical regions was common, occurring in 20 cases (37.0%). Information about the firing distance was available in 38 cases and in 26 cases (59.1%) the gunshot was made at “distance”.

In accidents one entrance wound was achieved in 6 cases (60.0%), being the abdomen the main region involved ( $n = 4$ , 40.0%), closely followed by head and/or neck with 3 cases (30.0%). Fifty percent of gunshots occurred at “near contact”. In both homicides and accidents were not recorded shots in contact.

Using the Chi-Square Test, it was possible to observe that the location of entrance wounds and firing distance were associated with the manner of death ( $p = 0.001$ ).

In the toxicological study, the presence of alcohol was positive in 19 cases of homicides with concentrations ranging from 0.13g/L to 2.57g/L, while in suicides concentrations ranged from 0.67g/L to 3.45g/L, in six positive cases.

## Discussion

Shotguns are used in a significant number of homicides and suicides, however, in literature the characterization of cases where this type of weapon is used is rare [13].

Between 2005 and 2010 the total number of deaths involving shotguns was similar in northern Portugal, with no pattern of evolution. In this period 105 cases were recorded in the northern region of Portugal (0.46/100000 inhabitants in north region, in 2010).

Of these cases, 51.4% ( $n = 54$ ) were homicides, 39.0% ( $n = 41$ ) suicides and 9.5% ( $n = 10$ ) accidents. In USA, a study showed the use of shotguns in 46.5% homicides, 52.4% suicides and 0.7% accidents [13]. Singh et al [16] reported 39.0% of homicides, 3% of suicides and 19% of accidents in a region of India. These results clearly show a variability in the use of this weapon depending on geographical aspects, legal rules and traditions of each country [9,16].

The use of firearms in violent deaths is associated predominantly with males, being men the main victims of shotguns [13]. In our study this pattern was observed, with 81.0% of male victims. This difference was smaller in the case of homicides with 35.2% of female victims. With regard to suicide cases, which are more fully described in the literature, in our study we obtained a ratio of male to female 40 (40/1), much higher compared to other studies [2]. This difference is explained by the existence of cultural patterns that associate men with more violent deaths [19]. In addition, men have higher suicidal tendencies, greater familiarity with the method and less concern with the disfiguration caused by the weapon, unlike women who seek to avoid methods that alter body image [19]. According to Kohlmeier et al [20], the handgun was the most common firearm chosen to commit suicide by females.

Our data shows no differences in the distribution of cases by age starting from age 25yo. Homicides with firearms tend to occur more in adults, being less frequent in young people and the elderly [14]. In our study the victims were mainly adults, the group [25-64yo] had 72.2% of cases. Miller et al

[21] revealed that in USA, most attempts of suicide occurred among people aged 15-24yo, but deaths were more frequent in adults older than 24 years. In our study there was only one case under 25 years, which shows that the use of these guns by young people is rare in Portugal. The same was found in homicides.

In accidents all victims were men. This predominance was also found in studies from others countries like USA, Denmark and Sweden [8]. Death in such cases occurs more often in young people, aged between [15-24yo] [8]. In our study the highest number of cases occurred with victims under 25 years old (40.0%) and two of these cases occurred when youths were playing with guns. This arouses attention to the need for greater caution of caregivers in accessing children/young people to arms, whether they are loaded or not [8].

The studies support the hypothesis that the use of shotguns in suicides, homicides and accidents is higher in rural areas compared to urban areas [18]. The data collected in our study regarding the victims' residence indicated greater proportion of victims in rural areas (66.7%). In these regions this type of weapon is easily acquired [16] and is traditionally transmitted between generations in the family [6], founding them in many houses hanged on the walls [16].

Farming was the main occupation (25.0%). This finding is consistent with data from other study [16]. It is known that this occupation has a high risk of suicide with firearms [21]. However, both the percentage of fatalities in rural areas as their occupation with agriculture may be undervalued. This could result from a decrease of inhabitants in rural areas where it is more frequent and also by the lack of information about the occupations in some cases, in particular previous occupations of retirees (36.6%).

Although cardiovascular diseases constitute the main pathology found on the victims, it is important to highlight the high number of suicides in which victims have diagnosed psychiatric disorders (47.8%). While there is no unanimity in the association between history of psychiatric illness and suicide, according to Cavanagh et al [22], psychiatric disease, especially diseases of humor, is present in 90% of victims who commit suicide.

The investigation of the motive/scenario is an essential component when analyzing violent deaths [23]. In homicides the problems in marital relationship (32.1%), discussions with neighbors (25.0%) and robberies (21.4%) were the main causes. These situations reflect the easy accessibility to obtain this type of weapon and its presence at home, making it a great tool to solve individuals' problems in a violent manner.

The data collected showed that in the case of suicides psychiatric diseases was the precipitating factor in 25.0% of cases. This shows that despite their diagnosis and respective treatment, it may not be effective or not fulfilled, so in these cases access to firearms should be actively restricted and the physician perform an important role in prevention [17]. It is also important to note that the presence of other disorders was found in 20% of cases. This scenario alert to the fact that physicians have also an important role in this type of situation [22]. They should understand how each individual reacts to disease, because even if their life may not be at risk, other factors such as loss of independence may be important triggers of this act [22].

With regard to accidents, shotguns are the type of weapons more susceptible to carelessness in handling and are used more often in risky situations, such as hunting [8]. In our study, the accidents

occurred mainly by the mishandling of the gun (40.0%) and hunting accidents (30.0%), which are the motives most frequently reported in other studies [8,16]

There is scientific evidence that indicates that people who consume alcohol or illicit drugs have an increased risk of committing acts of violence to others or to themselves, however, alcohol abuse is more strongly associated with violent deaths than illegal drugs [24]. In our study, regular consumption of alcohol was found in 53.6% of cases. This shows that the carrying of firearms by these individuals should be limited, having families and the physician an important role [25].

The severity and lethality of shotgun wounds depends on the number of pellets that enter the body, the organ hit by the pellets and the amount of tissue destroyed [16]. This fact is supported in our study because medical intervention occurred in only 16 cases (15.2%), showing that this weapon is very lethal.

Shotgun wounds are different depending on the type of pellets used, the distance of the shot and the region affected [26]. In accordance to other studies [13,16], our data showed that the most affected sites were head and/or neck (75.6%), followed by the chest (12.2%) and abdomen (12.2%).

Molina et al [13] revealed that the achievement of regions that host important organs such as the head and chest and even multiple locations occur more often in homicides, but there is no significant statistically evidence of a preferential anatomic site. Our study showed that entrance wounds were located on the head and/or neck and chest in 29.6%, and the involvement of various anatomical regions simultaneously occurred in 37.0% of cases. In suicides with firearm the head is generally the involved site [10]. In our study, most suicides were accomplished by shots fired in the head and/or neck (75.6%), which was consistent with previous studies [2,13]. In accidents, unlike other studies [8,16], the abdomen was the most affected site, followed by the head. However, in these cases there are no anatomical regions preferred and the firing distance is variable also too [13,16].

In shooting at short distances, the shotgun is the most tremendous and destructive of small guns [17]. The distance determination of the shots is performed by evaluating the presence or absence of the residues that leave the firing barrel [27]. In gunshots made at contact, these elements penetrate the wound [27]. In “near contact” and intermediate distance there is a deposition of discharge residues around the orifice or directly on clothing that are covering the impact area [27]. As the muzzle of the shotgun is moved farther from the body, tattooing disappears and the diameter of the circular wound of entrance increases in size and individual pellet holes may appear [17].

In homicides the most common firing distance was distant [13]. Our data showed the same results. In suicides “near contact” wounds were the most common, which can be explained by the absence of a device to pull the trigger, that is necessary to achieve longer shooting distances [15]. It was demonstrated a relationship between the location of entrance wounds and firing distance with the manner of death ( $p = 0.001$ ).

In the toxicology study, analysis of blood alcohol was performed more times, followed by analysis of illegal drugs and therapeutic drugs. In our study we revealed that 26.8% of the victims where a toxicological study was performed had alcohol in their blood. Karger et al [9] revealed that the blood alcohol analysis may be useful in establishing the manner of death, because the homicide victims had blood alcohol less frequently than the suicide victims. Our data showed that in homicides 19 cases were

positive, the concentration ranged from 0.13g/L to 2.57g/L, while in suicides concentration ranged from 0.67g/L to 3.45g/L, in six positive cases. It is known that men have a higher probability of being intoxicated before committing suicide [28].

### Limitations

The analysis of some variables were limited due to the retrospective nature of the study, therefore it was not possible to collect all data. Some information collected is based on social surveys, which does not allow us to have full credibility on the data and, moreover, in many cases there is no information at all.

### **Conclusion**

The determination of the manner of death (suicide, homicide or accidents) is one of the main objectives of the Forensic Medicine. The studies that seek to analyze series of cases are an important way to understand and find common denominators in each situation, leading forensic science to a preventive attitude of violent deaths.

There are many variables that contribute in a significant way for this diagnosis: investigation of the death scene and the circumstances preceding the death, victims' characteristics, gun characterization (type, number of shots and shooting distance) and the location of the entrance wound.

Information about the use of shotguns is scarce in the literature however, due to a significant use of these weapons in violent deaths and also a significant presence in the homes of Portuguese people, it is important to obtain a characterization of their use.

In our study, the use of shotguns occurred in 105 cases, which shows that despite restrictive laws in Portugal the use of these weapons is not so scarce. Our data shows that its use is predominant in homicides and suicides and men are the main victims. They are used mainly in rural areas and by farmers. Young people seem to assume a significant role in accidents.

The head and/or neck were the most involved regions however, there are variations depending on the manner of death that can help to point it out. Firing distance seems to have a similar role.

The motives are also closely related to the manner of death: in homicides, marital discussions or with neighbors were the main causes; in suicides diseases are the precipitating reasons found in more cases and in accidents we have mishandling of the gun and hunting accidents.

Therefore, it is important to understand the real existence of these weapons, because knowing this information can induce a change in mentalities to a more restrictive law that could reduce the number of violent deaths by firearms.

### **Integrity of research and reporting**

#### Ethical standards

The authors developed a retrospective study and were not carried out any intervention.

#### Conflict of interest

The authors declare that they have no conflict of interest.

## References

1. WHO (2001) Small Arms and Global Health. [http://www.who.int/violence\\_injury\\_prevention/index.html](http://www.who.int/violence_injury_prevention/index.html). Accessed 12 November 2012
2. Asirdizer M, Canturk G, Canturk N, Yavuz MS, Sari H (2010) Analyses of suicidal deaths with shotguns in Istanbul, 1998-2007. *Ulus Travma Acil Cerrahi Derg* 16 (1):47-53
3. Karch DL, Logan J, McDaniel D, Parks S, Patel N (2012) Surveillance for violent deaths--National Violent Death Reporting System, 16 states, 2009. *MMWR Surveill Summ* 61 (6):1-43
4. Justiça Md (2005-2010) Estatísticas da Justiça - Armas Utilizadas em Crimes Registrados [www.dgpi.mj.pt](http://www.dgpi.mj.pt). Accessed 19 November 2012
5. Dahlberg LL, Ikeda RM, Kresnow MJ (2004) Guns in the home and risk of a violent death in the home: findings from a national study. *Am J Epidemiol* 160 (10):929-936.
6. Cardoso J, Gonçalves F, Campos J (2011) Armas de Fogo de Cano de Alma Lisa - Contribuição para análise das características do disparo. Dissertation.
7. CES (2010) Violência e armas ligeiras: um retrato Português. <http://www.ces.uc.pt/iframe/nucleos/nep/pdfs/RelatorioSeminarioArmas.pdf>. Accessed 19 November 2012
8. Karger B, Billeb E, Koops E (2002) Accidental firearm fatalities. Forensic and preventive implications. *Int J Legal Med* 116 (6):350-353.
9. Karger B, Billeb E, Koops E, Brinkmann B (2002) Autopsy features relevant for discrimination between suicidal and homicidal gunshot injuries. *Int J Legal Med* 116 (5):273-278.
10. Desinan L, Mazzolo GM (2005) Gunshot fatalities: suicide, homicide or accident? A series of 48 cases. *Forensic Sci Int* 147 Suppl:S37-40.
11. Rainio J, Sajantila A (2005) Fatal gunshot wounds between 1995 and 2001 in a highly populated region in Finland. *Am J Forensic Med Pathol* 26 (1):70-77
12. Balci Y, Canogullari G, Ulupinar E (2007) Characterization of the gunshot suicides. *J Forensic Leg Med* 14 (4):203-208.

13. Molina DK, Wood LE, DiMaio VJ (2007) Shotgun wounds: a review of range and location as pertaining to manner of death. *Am J Forensic Med Pathol* 28 (2):99-102.
14. Solarino B, Nicoletti EM, Di Vella G (2007) Fatal firearm wounds: a retrospective study in Bari (Italy) between 1988 and 2003. *Forensic Sci Int* 168 (2-3):95-101.
15. Grandmaison GL, Fermanian C, Aegerter P, Durigon M (2008) Influence of ballistic and autopsy parameters on the manner of death in case of long firearms fatalities. *Forensic Sci Int* 177 (2-3):207-213.
16. Singh BP, Singh RP (2005) Shotgun shooting in northern India--a review (1980-1999). *Forensic Sci Int* 150 (1):103-111.
17. DiMaio V (1999) *Gunshot Wounds - Practical aspects of firearms, ballistics, and forensic techniques*. CRC Press, New York
18. Dresang LT (2001) Gun deaths in rural and urban settings: recommendations for prevention. *J Am Board Fam Pract* 14 (2):107-115
19. Hawton K (2000) Sex and suicide. Gender differences in suicidal behaviour. *Br J Psychiatry* 177:484-485
20. Kohlmeier RE, McMahan CA, DiMaio VJ (2001) Suicide by firearms: a 15-year experience. *Am J Forensic Med Pathol* 22 (4):337-340
21. Miller M, Azrael D, Hemenway D (2004) The epidemiology of case fatality rates for suicide in the northeast. *Ann Emerg Med* 43 (6):723-730.
22. Cavanagh JT, Carson AJ, Sharpe M, Lawrie SM (2003) Psychological autopsy studies of suicide: a systematic review. *Psychol Med* 33 (3):395-405
23. Coelho L, Ribeiro T, Dias R, Santos A, Magalhaes T (2010) Elder homicide in the north of Portugal. *J Forensic Leg Med* 17 (7):383-387.
24. Hawton K, van Heeringen K (2009) Suicide. *Lancet* 373 (9672):1372-1381.
25. Webster DW, Vernick JS (2009) Keeping firearms from drug and alcohol abusers. *Inj Prev* 15 (6):425-427.
26. Abe M, Motani-Saitoh H, Sato Y, Kiuchi M (2002) A fatal case of shotgun injury caused by one pellet. *Leg Med (Tokyo)* 4 (2):131-133

27. Denton JS, Segovia A, Filkins JA (2006) Practical pathology of gunshot wounds. *Arch Pathol Lab Med* 130 (9):1283-1289.
28. Kaplan MS, McFarland BH, Huguet N (2009) Characteristics of adult male and female firearm suicide decedents: findings from the National Violent Death Reporting System. *Inj Prev* 15 (5):322-327.

## Tables

Table 1 – Social and demographic characteristics of the victims.

|                       |            | <i>n</i> | %    | Total n (%) |
|-----------------------|------------|----------|------|-------------|
| <b>Gender</b>         | Male       | 85       | 81.0 | 105 (100)   |
|                       | Female     | 20       | 19.0 |             |
| <b>Age</b>            | [15-24yo]  | 6        | 5.7  | 105 (100.0) |
|                       | [25-34yo]  | 17       | 16.2 |             |
|                       | [35-44yo]  | 15       | 14.3 |             |
|                       | [45-54yo]  | 18       | 17.1 |             |
|                       | [55-64yo]  | 21       | 20.0 |             |
|                       | [65-74yo]  | 10       | 9.5  |             |
|                       | [+75yo]    | 18       | 17.1 |             |
| <b>Marital Status</b> | Married    | 66       | 63.5 | 104 (99.0)  |
|                       | Single     | 20       | 19.2 |             |
|                       | Divorced   | 13       | 12.5 |             |
|                       | Widowed    | 5        | 4.8  |             |
| <b>Job</b>            | Employed   | 46       | 50.0 | 92 (87.6)   |
|                       | Retired    | 34       | 37.0 |             |
|                       | Unemployed | 12       | 13.0 |             |
| <b>Residence</b>      | Rural      | 70       | 66.7 | 105 (100.0) |
|                       | Urban      | 35       | 33.3 |             |

Table 2 – Distribution of cases over the 6-years.

|             |      | <i>n</i> | %    | Total n (%) |
|-------------|------|----------|------|-------------|
| <b>Year</b> | 2005 | 21       | 20.0 | 105 (100.0) |
|             | 2006 | 15       | 14.3 |             |
|             | 2007 | 15       | 14.3 |             |
|             | 2008 | 18       | 17.1 |             |
|             | 2009 | 21       | 20.0 |             |
|             | 2010 | 15       | 14.3 |             |

Table 3 – Place of death

|                  | <i>n</i> | %    | Total n (%) |
|------------------|----------|------|-------------|
| Home             | 59       | 56.7 | 104 (99.0)  |
| Public Place     | 36       | 34.6 |             |
| Work's Place     | 5        | 4.8  |             |
| Neighbors' house | 3        | 2.9  |             |
| Families house   | 1        | 1.0  |             |



Table 4 – Forensic findings

|                                |                          | <i>n</i> | %    | Total n (%) |
|--------------------------------|--------------------------|----------|------|-------------|
| <b>Entrance wound location</b> | Head + Neck              | 50       | 47.6 |             |
|                                | Chest                    | 22       | 21.0 |             |
|                                | Abdomen                  | 10       | 9.5  |             |
|                                | Back                     | 2        | 1.9  |             |
|                                | Limbs                    | -        | -    |             |
|                                | Multiple wound locations | 20       | 37.0 | 105 (100.0) |
| <b>Firing distance</b>         | Contact                  | 4        | 3.8  |             |
|                                | “Near Contact”           | 28       | 26.7 |             |
|                                | Intermediate distance    | 11       | 10.5 |             |
|                                | Distant                  | 29       | 27.6 | 72 (68.6)   |

Table 5 – Social and demographic characteristics of the victims according to the manner of death

|                       |            | <b>Suicide (n,%)</b> |       | <b>Homicide (n,%)</b> |       | <b>Accident (n,%)</b> |       |
|-----------------------|------------|----------------------|-------|-----------------------|-------|-----------------------|-------|
| <b>Gender</b>         | Male       | 40                   | 97.6  | 35                    | 64.8  | 10                    | 100.0 |
|                       | Female     | 1                    | 2.4   | 19                    | 35.2  | -                     | -     |
|                       | Total      | 41                   | 100.0 | 54                    | 100.0 | 10                    | 100.0 |
| <b>Age</b>            | [15-24yo]  | 1                    | 2.4   | 1                     | 1.9   | 4                     | 40.0  |
|                       | [25-34yo]  | 5                    | 12.2  | 12                    | 22.2  | 1                     | 10.0  |
|                       | [35-44yo]  | 7                    | 17.1  | 8                     | 14.8  | -                     | -     |
|                       | [45-54yo]  | 5                    | 12.2  | 12                    | 22.2  | 1                     | 10.0  |
|                       | [55-64yo]  | 10                   | 24.4  | 7                     | 13.0  | 3                     | 30.0  |
|                       | [65-74yo]  | 5                    | 12.2  | 5                     | 9.3   | -                     | -     |
|                       | [+75yo]    | 8                    | 19.5  | 9                     | 16.7  | 1                     | 10.0  |
|                       | Total      | 41                   | 100.0 | 54                    | 100.0 | 10                    | 100.0 |
| <b>Marital Status</b> | Married    | 29                   | 70.7  | 30                    | 56.6  | 7                     | 70.0  |
|                       | Single     | 5                    | 12.2  | 12                    | 22.6  | 3                     | 30.0  |
|                       | Divorced   | 5                    | 12.2  | 8                     | 15.1  | -                     | -     |
|                       | Widowed    | 2                    | 4.9   | 3                     | 5.7   | -                     | -     |
|                       | Total      | 41                   | 100.0 | 54                    | 100.0 | 10                    | 100.0 |
| <b>Job</b>            | Employed   | 16                   | 44.4  | 22                    | 47.8  | 8                     | 80.0  |
|                       | Retired    | 17                   | 47.2  | 15                    | 32.6  | 2                     | 20.0  |
|                       | Unemployed | 3                    | 8.3   | 9                     | 19.6  | -                     | -     |
|                       | Total      | 36                   | 100.0 | 46                    | 100.0 | 10                    | 100.0 |
| <b>Residence</b>      | Rural      | 27                   | 65.9  | 36                    | 66.7  | 7                     | 70.0  |
|                       | Urban      | 14                   | 34.1  | 18                    | 33.3  | 3                     | 30.0  |
|                       | Total      | 41                   | 100.0 | 54                    | 100.0 | 10                    | 100.0 |

Table 6 – Place of event according to the manner of death.

|                  | Suicide (n,%) |      | Homicide (n,%) |      | Accident (n,%) |      |
|------------------|---------------|------|----------------|------|----------------|------|
| Home             | 37            | 90.2 | 20             | 37.0 | 3              | 30.0 |
| Public Place     | 4             | 9.8  | 26             | 48.1 | 6              | 60.0 |
| Work's Place     | -             | -    | 5              | 9.3  | -              | -    |
| Neighbors' house | -             | -    | 3              | 5.6  | -              | -    |
| Families house   | -             | -    | -              | -    | 1              | 10.0 |

Table 7 – Life events in relation to the manner of death.

|                 |                                      | n | %    | Total (n, %) |
|-----------------|--------------------------------------|---|------|--------------|
| <b>Suicide</b>  | Psychiatric disease                  | 5 | 29.5 |              |
|                 | Life-threatening illness             | 4 | 23.5 |              |
|                 | Job problem                          | 3 | 17.6 |              |
|                 | Problems in the marital relationship | 2 | 11.8 |              |
|                 | Terminal illness                     | 1 | 5.9  |              |
|                 | Financial problem                    | 1 | 5.9  |              |
|                 | Recent death of relative             | 1 | 5.9  | 17 (41.5)    |
| <b>Homicide</b> | Problems in the marital relationship | 9 | 34.6 |              |
|                 | Discussion with neighbor             | 7 | 26.9 |              |
|                 | Robbery                              | 6 | 23.1 |              |
|                 | Job problem                          | 2 | 7.7  |              |
|                 | Discussion with family member        | 2 | 7.7  | 26 (48.1)    |
| <b>Accident</b> | Mishandling of the gun               | 4 | 40.0 |              |
|                 | Hunting accident                     | 3 | 30.0 |              |
|                 | Playing with gun                     | 2 | 20.0 |              |
|                 | Drop with gun                        | 1 | 10.0 | 10 (100.0)   |

Table 8 – Forensic findings related to the manner of death.

|                 |   | Suicide (n,%) |      | Homicide (n,%) |      | Accident (n,%) |      |
|-----------------|---|---------------|------|----------------|------|----------------|------|
| <b>Number</b>   | 1 | 33            | 80.5 | 22             | 40.7 | 5              | 50.0 |
| <b>entrance</b> | 2 | -             | -    | 4              | 7.4  | 1              | 10.0 |
| <b>wounds</b>   | 3 | -             | -    | 1              | 1.9  | -              | -    |

|                        |                          |    |       |    |       |    |       |
|------------------------|--------------------------|----|-------|----|-------|----|-------|
|                        | Multiple                 | -  | -     | 26 | 48.1  | 2  | 20.0  |
|                        | Undetermined             | 8  | 19.5  | 1  | 1.9   | 2  | 20.0  |
|                        | Total                    | 41 | 100.0 | 54 | 100.0 | 10 | 100.0 |
| <b>Location</b>        | Head + Neck              | 31 | 75,6  | 16 | 29,6  | 3  | 30,0  |
|                        | Chest                    | 5  | 12,2  | 16 | 29,6  | 1  | 10,0  |
|                        | Abdomen                  | 5  | 12,2  | 1  | 1,9   | 4  | 40,0  |
|                        | Back                     | -  | -     | 1  | 1,9   | 1  | 10,0  |
|                        | Limbs                    | -  | -     | -  | -     | -  | -     |
|                        | Multiple wound locations | -  | -     | 20 | 37,0  | 1  | 10,0  |
|                        | Total                    | 41 | 100.0 | 54 | 100.0 | 10 | 100.0 |
| <b>Firing distance</b> | Contact                  | 4  | 21.1  | -  | -     | -  | -     |
|                        | Near Contact             | 15 | 78.9  | 8  | 18.2  | 5  | 55.6  |
|                        | Intermediate distance    | -  | -     | 10 | 22.7  | 1  | 11.1  |
|                        | Distant                  | -  | -     | 26 | 59.1  | 3  | 33.3  |
|                        | Total                    | 19 | 100.0 | 44 | 100.0 | 9  | 100.0 |

## **Annexes**

### **Annex 1: Instructions for Authors**

## **GUIDELINES FOR PUBLISHING POPULATION DATA**

In 1997 Prof. Bernd Brinkmann formulated guidelines for the submission of manuscripts on short tandem repeat (STR) population data (Brinkmann 1997). These earlier guidelines have now been extended to include haploid DNA markers, i.e. mitochondrial DNA (mtDNA) and Y-chromosomal polymorphisms.

For specific information, see the Short Communication “Publication of population data of linearly inherited DNA markers in the International Journal of Legal Medicine” (Parson and Roewer 2010; DOI 10.1007/s00414-010-0492-y) published online in Int J Legal Med in July 2010.

All forensic population genetics papers should always contain information on the description of the population, ethical requirements and quality control. For mtDNA papers, previous acceptance of the dataset in EMPOP is required; for YSTR and YSNP data, previous inclusion of the data in the YSTR/YSNP database is required.

- EMPOP database
- YSTR/YSNP database

## **MANUSCRIPT SUBMISSION**

### **Manuscript Submission**

Submission of a manuscript implies: that the work described has not been published before; that it is not under consideration for publication anywhere else; that its publication has been approved by all co-authors, if any, as well as by the responsible authorities – tacitly or explicitly – at the institute where the work has been carried out.

The publisher will not be held legally responsible should there be any claims for compensation.

### **Permissions**

Authors wishing to include figures, tables, or text passages that have already been published elsewhere are required to obtain permission from the copyright owner(s) for both the print and online format and to include evidence that such permission has been granted when submitting their papers. Any material received without such evidence will be assumed to originate from the authors.

### **Online Submission**

Authors should submit their manuscripts online. Electronic submission substantially reduces the editorial processing and reviewing times and shortens overall publication times. Please follow the hyperlink “Submit online” on the right and upload all of your manuscript files following the instructions given on the screen.

## **TITLE PAGE**

### **Title Page**

The title page should include:

- The name(s) of the author(s)
- A concise and informative title
- The affiliation(s) and address(es) of the author(s)
- The e-mail address, telephone and fax numbers of the corresponding author

### **Abstract**

Please provide an abstract of 150 to 250 words. The abstract should not contain any undefined abbreviations or unspecified references.

### **Keywords**

Please provide 4 to 6 keywords which can be used for indexing purposes.

## **TEXT**

### **Text Formatting**

Manuscripts should be submitted in Word.

- Use a normal, plain font (e.g., 10-point Times Roman) for text.
- Use italics for emphasis.
- Use the automatic page numbering function to number the pages.
- Do not use field functions.
- Use tab stops or other commands for indents, not the space bar. Use the table function, not spreadsheets, to make tables.
- Use the equation editor or MathType for equations.  
Note: If you use Word 2007, do not create the equations with the default equation editor but use the Microsoft equation editor or MathType instead.
- Save your file in doc format. Do not submit docx files.  
Word template (zip, 154 kB)

Manuscripts with mathematical content can also be submitted in LaTeX.

LaTeX macro package (zip, 182 kB)

### **Headings**

Please use no more than three levels of displayed headings.

### **Abbreviations**

Abbreviations should be defined at first mention and used consistently thereafter.

## Footnotes

Footnotes can be used to give additional information, which may include the citation of a reference included in the reference list. They should not consist solely of a reference citation, and they should never include the bibliographic details of a reference. They should also not contain any figures or tables.

Footnotes to the text are numbered consecutively; those to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data). Footnotes to the title or the authors of the article are not given reference symbols.

Always use footnotes instead of endnotes.

## Acknowledgments

Acknowledgments of people, grants, funds, etc. should be placed in a separate section before the reference list. The names of funding organizations should be written in full.

## REFERENCES

### Citation

Reference citations in the text should be identified by numbers in square brackets. Some examples:

1. Negotiation research spans many disciplines [3].
2. This result was later contradicted by Becker and Seligman [5].
3. This effect has been widely studied [1-3, 7].

### Reference list

The list of references should only include works that are cited in the text and that have been published or accepted for publication. Personal communications and unpublished works should only be mentioned in the text.

Do not use footnotes or endnotes as a substitute for a reference list.

The entries in the list should be numbered consecutively.

- Journal article

Gamelin FX, Baquet G, Berthoin S, Thevenet D, Nourry C, Nottin S, Bosquet L (2009) Effect of high intensity intermittent training on heart rate variability in prepubescent children. *Eur J Appl Physiol* 105:731-738. doi: 10.1007/s00421-008-0955-8

Ideally, the names of all authors should be provided, but the usage of “et al” in long author lists will also be accepted:

Smith J, Jones M Jr, Houghton L et al (1999) Future of health insurance. *N Engl J Med* 341:325–329

- Article by DOI

Slifka MK, Whitton JL (2000) Clinical implications of dysregulated cytokine production. *J Mol Med*. doi: 10.1007/s001090000086

- Book  
South J, Blass B (2001) The future of modern genomics. Blackwell, London
- Book chapter  
Brown B, Aaron M (2001) The politics of nature. In: Smith J (ed) The rise of modern genomics, 3<sup>rd</sup> edn. Wiley, New York, pp 230-257
- Online document  
Cartwright J (2007) Big stars have weather too. IOP Publishing PhysicsWeb. <http://physicsweb.org/articles/news/11/6/16/1>. Accessed 26 June 2007
- Dissertation  
Trent JW (1975) Experimental acute renal failure. Dissertation, University of California

Always use the standard abbreviation of a journal's name according to the ISSN List of Title Word Abbreviations, see [www.issn.org/2-22661-LTWA-online.php](http://www.issn.org/2-22661-LTWA-online.php)

For authors using EndNote, Springer provides an output style that supports the formatting of in-text citations and reference list.

EndNote style (zip, 2 kB)

## **TABLES**

- All tables are to be numbered using Arabic numerals.
- Tables should always be cited in text in consecutive numerical order.
- For each table, please supply a table caption (title) explaining the components of the table.
- Identify any previously published material by giving the original source in the form of a reference at the end of the table caption.
- Footnotes to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data) and included beneath the table body.

## **ARTWORK**


For the best quality final product, it is highly recommended that you submit all of your artwork – photographs, line drawings, etc. – in an electronic format. Your art will then be produced to the highest standards with the greatest accuracy to detail. The published work will directly reflect the quality of the artwork provided.

## **Electronic Figure Submission**

- Supply all figures electronically.
- Indicate what graphics program was used to create the artwork.
- For vector graphics, the preferred format is EPS; for halftones, please use TIFF format. MS Office files are also acceptable.


- Vector graphics containing fonts must have the fonts embedded in the files.
- Name your figure files with "Fig" and the figure number, e.g., Fig1.eps.

### Line Art



- Definition: Black and white graphic with no shading.
- Do not use faint lines and/or lettering and check that all lines and lettering within the figures are legible at final size.
- All lines should be at least 0.1 mm (0.3 pt) wide.
- Scanned line drawings and line drawings in bitmap format should have a minimum resolution of 1200 dpi.
- Vector graphics containing fonts must have the fonts embedded in the files.

### Halftone Art




- Definition: Photographs, drawings, or paintings with fine shading, etc.



- If any magnification is used in the photographs, indicate this by using scale bars within the figures themselves.
- Halftones should have a minimum resolution of 300 dpi.

### Combination Art



- Definition: a combination of halftone and line art, e.g., halftones containing line drawing, extensive lettering, color diagrams, etc.
- Combination artwork should have a minimum resolution of 600 dpi.

### Color Art

- Color art is free of charge for online publication.
- If black and white will be shown in the print version, make sure that the main information will still be visible. Many colors are not distinguishable from one another when converted to black and white. A simple way to check this is to make a xerographic copy to see if the necessary distinctions between the different colors are still apparent.
- If the figures will be printed in black and white, do not refer to color in the captions.
- Color illustrations should be submitted as RGB (8 bits per channel).

### Figure Lettering

- To add lettering, it is best to use Helvetica or Arial (sans serif fonts).
- Keep lettering consistently sized throughout your final-sized artwork, usually about 2–3 mm (8–12 pt).

- Variance of type size within an illustration should be minimal, e.g., do not use 8-pt type on an axis and 20-pt type for the axis label.
- Avoid effects such as shading, outline letters, etc.
- Do not include titles or captions within your illustrations.

### **Figure Numbering**

- All figures are to be numbered using Arabic numerals.
- Figures should always be cited in text in consecutive numerical order.
- Figure parts should be denoted by lowercase letters (a, b, c, etc.).
- If an appendix appears in your article and it contains one or more figures, continue the consecutive numbering of the main text. Do not number the appendix figures, "A1, A2, A3, etc." Figures in online appendices (Electronic Supplementary Material) should, however, be numbered separately.

### **Figure Captions**

- Each figure should have a concise caption describing accurately what the figure depicts. Include the captions in the text file of the manuscript, not in the figure file.
- Figure captions begin with the term Fig. in bold type, followed by the figure number, also in bold type.
- No punctuation is to be included after the number, nor is any punctuation to be placed at the end of the caption.
- Identify all elements found in the figure in the figure caption; and use boxes, circles, etc., as coordinate points in graphs.
- Identify previously published material by giving the original source in the form of a reference citation at the end of the figure caption.

### **Figure Placement and Size**

- When preparing your figures, size figures to fit in the column width.
- For most journals the figures should be 39 mm, 84 mm, 129 mm, or 174 mm wide and not higher than 234 mm.
- For books and book-sized journals, the figures should be 80 mm or 122 mm wide and not higher than 198 mm.

### **Permissions**

If you include figures that have already been published elsewhere, you must obtain permission from the copyright owner(s) for both the print and online format. Please be aware that some publishers do not grant electronic rights for free and that Springer will not be able to refund any costs that may have occurred to receive these permissions. In such cases, material from other sources should be used.

### **Accessibility**

In order to give people of all abilities and disabilities access to the content of your figures, please make sure that

- All figures have descriptive captions (blind users could then use a text-to-speech software or a text-to-Braille hardware)
- Patterns are used instead of or in addition to colors for conveying information (color-blind users would then be able to distinguish the visual elements)
- Any figure lettering has a contrast ratio of at least 4.5:1

## **ELECTRONIC SUPPLEMENTARY MATERIAL**

Springer accepts electronic multimedia files (animations, movies, audio, etc.) and other supplementary files to be published online along with an article or a book chapter. This feature can add dimension to the author's article, as certain information cannot be printed or is more convenient in electronic form.

### **Submission**

- Supply all supplementary material in standard file formats.
- Please include in each file the following information: article title, journal name, author names; affiliation and e-mail address of the corresponding author.
- To accommodate user downloads, please keep in mind that larger-sized files may require very long download times and that some users may experience other problems during downloading.

### **Audio, Video, and Animations**

- Always use MPEG-1 (.mpg) format.

### **Text and Presentations**

- Submit your material in PDF format; .doc or .ppt files are not suitable for long-term viability.
- A collection of figures may also be combined in a PDF file.

### **Spreadsheets**

- Spreadsheets should be converted to PDF if no interaction with the data is intended.
- If the readers should be encouraged to make their own calculations, spreadsheets should be submitted as .xls files (MS Excel).

### **Specialized Formats**

- Specialized format such as .pdb (chemical), .vrl (VRML), .nb (Mathematica notebook), and .tex can also be supplied.

### **Collecting Multiple Files**

- It is possible to collect multiple files in a .zip or .gz file.

### **Numbering**

- If supplying any supplementary material, the text must make specific mention of the material as a citation, similar to that of figures and tables.
- Refer to the supplementary files as “Online Resource”, e.g., "... as shown in the animation (Online Resource 3)", "... additional data are given in Online Resource 4”.
- Name the files consecutively, e.g. “ESM\_3.mpg”, “ESM\_4.pdf”.

### **Captions**

- For each supplementary material, please supply a concise caption describing the content of the file.

### **Processing of supplementary files**

- Electronic supplementary material will be published as received from the author without any conversion, editing, or reformatting.

### **Accessibility**

- In order to give people of all abilities and disabilities access to the content of your supplementary files, please make sure that
- The manuscript contains a descriptive caption for each supplementary material
- Video files do not contain anything that flashes more than three times per second (so that users prone to seizures caused by such effects are not put at risk)

## **INTEGRITY OF RESEARCH AND REPORTING**

### **Ethical standards**

Manuscripts submitted for publication must contain a declaration that the experiments comply with the current laws of the country in which they were performed. Please include this note in a separate section before the reference list.

### **Conflict of interest**

Authors must indicate whether or not they have a financial relationship with the organization that sponsored the research. This note should be added in a separate section before the reference list.

If no conflict exists, authors should state: The authors declare that they have no conflict of interest.

### **Animal Welfare**

If applicable, the author attests that experiments conducted on animal subjects complied with all applicable laws, regulations, and standards in the country where the studies were performed.

In general, it is expected that animal experimentation published in the International Journal of Legal Medicine complies with prevailing standards in either the European Union or the United States.

European Union standards

United States standards

## **AFTER ACCEPTANCE**

Upon acceptance of your article you will receive a link to the special Author Query Application at Springer's web page where you can sign the Copyright Transfer Statement online and indicate whether you wish to order OpenChoice, offprints, or printing of figures in color.

Once the Author Query Application has been completed, your article will be processed and you will receive the proofs.

### **Open Choice**

In addition to the normal publication process (whereby an article is submitted to the journal and access to that article is granted to customers who have purchased a subscription), Springer provides an alternative publishing option: Springer Open Choice. A Springer Open Choice article receives all the benefits of a regular subscription based article, but in addition is made available publicly through Springer's online platform SpringerLink. We regret that Springer Open Choice cannot be ordered for published articles.

Springer Open Choice

### **Copyright transfer**

Authors will be asked to transfer copyright of the article to the Publisher (or grant the Publisher exclusive publication and dissemination rights). This will ensure the widest possible protection and dissemination of information under copyright laws.

Open Choice articles do not require transfer of copyright as the copyright remains with the author. In opting for open access, they agree to the Springer Open Choice Licence.

### **Offprints**

Offprints can be ordered by the corresponding author.

### **Color illustrations**

Online publication of color illustrations is free of charge. For color in the print version, authors will be expected to make a contribution towards the extra costs.

### **Proof reading**

The purpose of the proof is to check for typesetting or conversion errors and the completeness and accuracy of the text, tables and figures. Substantial changes in content, e.g., new results, corrected values, title and authorship, are not allowed without the approval of the Editor.

After online publication, further changes can only be made in the form of an Erratum, which will be hyperlinked to the article.

### **Online First**

The article will be published online after receipt of the corrected proofs. This is the official first publication citable with the DOI. After release of the printed version, the paper can also be cited by issue and page numbers.

#### **SCIENTIFIC STYLE**

- Please always use internationally accepted signs and symbols for units, SI units.
- Genus and species names should be in italics.